

The septin cortex at the yeast mother–bud neck

Amy S Gladfelter*, John R Pringle† and Daniel J Lew‡

A specialized cortical domain is organized by the septins at the necks of budding yeast cells. Recent findings suggest that this domain serves as a diffusion barrier and also as a local cell-shape sensor. We review these findings along with what is known about the organization of the septin cortex and its regulation during the cell cycle.

Addresses

*Department of Molecular Microbiology, Biozentrum, University of Basel, Klingelbergstrasse 70, CH-4056 Basel, Switzerland

†Department of Biology and Program in Molecular Biology and Biotechnology, University of North Carolina, Chapel Hill, North Carolina 27599-3280, USA

‡Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, North Carolina 27710, USA; e-mail: daniel.lew@duke.edu

Current Opinion in Microbiology 2001, 4:681–689

1369-5274/01/\$ – see front matter

© 2001 Elsevier Science Ltd. All rights reserved.

Abbreviation

GFP green fluorescent protein


Introduction

The ability to specialize specific domains of the cell cortex is crucial to the function of almost all cell types. Notable examples include the distinct apical and basolateral membrane domains of epithelial cells, the pre- and post-synaptic membranes of neurons, the leading edges of migrating cells, and the cleavage furrows of dividing cells. In the budding yeast *Saccharomyces cerevisiae*, a specialized cortical domain is established that separates the bud, which will grow into a new daughter cell, from the pre-existing mother cell. The neck between mother and bud will become the site of cytokinesis and septum formation at the end of the cell cycle.

Throughout the process of bud formation, the cell cortex at the neck is clearly differentiated from the cortex at other parts of the cell, both inside and outside the plasma membrane (Figure 1). Outside, a chitin-rich ring (the eventual ‘bud scar’) forms in the cell wall on the mother side of the neck [1–4]. Inside, a family of evolutionarily conserved proteins called septins [5] form a highly ordered cortical domain that can be seen as a set of ~20 evenly-spaced striations spanning the neck by electron microscopy [6,7]. Although the neck is unique to budding yeasts, septins are present widely, if not ubiquitously, in animal cells, in which they may also serve to organize cortical domains, including regions concerned with cytokinesis [8–12] or with regulated secretion [13,14].

The septin-containing cortex in yeast is thought to act as a scaffold, recruiting many other proteins to the neck, where they can perform their varied functions (reviewed

Figure 1



The yeast mother–bud neck. (a) Scanning electron micrographs of haploid yeast cells. Note the bud scars from previous cell divisions. Adapted, with permission, from Chant and Pringle [59]. (b) Thin-section transmission electron micrograph of a small-budded cell. Note the alternating dark and light striations (arrowed) that underlie the plasma membrane. These striations disappear in septin-mutant strains at restrictive temperature. Adapted, with permission, from Byers and Goetsch [6]. The bar in (b) represents 0.1 μm.

by Longtine *et al.* [5]). Well-established functions for the septin cortex include synthesis of the chitin ring and septum, bud-site selection and cytokinesis [15–17]. The ability of the septins to recruit chitin synthases, bud-site-selection landmarks, actomyosin ring components and other proteins to the neck is perhaps sufficient to account for their role in these processes. However, recent findings have documented novel roles for the septins that appear to entail more than simply recruiting other proteins to the neck. These studies highlight longstanding questions regarding how the septin cortex is organized and how it is regulated during the cell cycle. In this review, we discuss these recent findings with a focus on the regulation of septin organization and its implications for septin function.

The septin cortex as diffusion barrier

In a beautiful set of experiments, Takizawa and colleagues [18••] demonstrated that the septin cortex serves as a diffusion barrier for integral membrane proteins. They first identified an mRNA that was transported to the bud tip by the actin cytoskeleton and then found that the encoded transmembrane protein, Ist2p, was localized exclusively to the bud plasma membrane. They suggested that translation, translocation and transport of Ist2p to the plasma membrane all occurred within the bud and, furthermore, that once at the membrane, Ist2p could not diffuse into the mother

portion of the cell. Using GFP-tagged Ist2p, they monitored fluorescence recovery after photobleaching of a spot on the bud membrane and concluded that Ist2p was freely diffusible within the plane of the membrane, indicating that the restriction to the bud was due to a diffusion barrier at the neck. In septin-mutant cells, this barrier was eliminated and Ist2p spread throughout the mother, as well as the bud membrane. The septin cortex's role as a diffusion barrier may explain why it is so highly ordered — perhaps the striations observed by electron microscopy correspond to the physical barriers to diffusion through the neck.

This barrier role of the septins invites comparison to tight junctions in animal cells, which exhibit clear 'ridges' of transmembrane proteins when examined by freeze-fracture electron microscopy. Thus far, no analogous structures have been described at the yeast neck. Given that the septins themselves seem to be cytoplasmic proteins that are only peripherally associated with the plasma membrane, it seems possible that the barrier in yeast only blocks the migration of membrane-associated molecules with sizable cytoplasmic domains and not that of all membrane components, including lipids. In this regard, it is interesting that another recent study suggested that the neck serves as a diffusion barrier for proteins that are peripherally associated with the plasma membrane [19^{*}]. However, this same study found that cortical actin patches (large mobile disc-shaped structures containing F-actin that are associated with the cytoplasmic face of the plasma membrane [20,21]) were able to migrate from the bud side of the neck to the mother. This astonishing feat would seem to imply that the barrier is highly selective and perhaps able to open to let the patches through. Alternatively, the patches may transiently abandon the plasma membrane to travel through the neck interior. Support for the latter possibility is provided by recent findings on actin patches in fission yeast, which do appear to make excursions away from the membrane into the cell interior [22]. The degree to which the septin cortex forms an effective barrier to the diffusion of various cellular constituents is clearly an issue worthy of further investigation. It seems likely that an understanding of this issue will require elucidation of the higher-order organization of the septins and associated proteins within this domain.

Regulators of septin organization

How are the septins organized within the neck cortex? This fundamental question remains largely unanswered. When Byers and Goetsch first examined the neck cortex by electron microscopy [6,7], they hypothesized that the striations represented helical filaments that wrapped around the neck (Figure 2a). Consistent with this proposal, overproduction of certain proteins can induce cells to elaborate extensive rings and spirals of septins into the bud [23^{*},24]. Subsequent studies demonstrated that purified septin complexes could polymerize to form filaments *in vitro* [25,26], and this capability seems likely to be important for the higher-order organization of the septin cortex *in vivo* (but see Field and


Kellogg [27] for a stimulating discussion on this point). These studies also led Field *et al.* [25] to propose an alternative model for septin organization in which polymerized septin filaments were oriented along the mother–bud axis and associated laterally to form a collar around the neck (Figure 2a). Support for this model has come from observations on mutant strains lacking factors important for septin organization. The first such factor to be identified was the protein kinase Gin4p [28], which is co-localized with septins at the neck (Figure 4). Mutant cells lacking Gin4p frequently display aberrant septin staining, in which several thick septin-containing bars are oriented along the mother–bud axis (Figure 2b). Similar bars have also been observed more recently in cells lacking the Gin4p-binding protein Nap1p or the Cdc42p-activated kinase Cla4p [29^{**}]. An attractive interpretation of the mutant phenotype is that Gin4p, Nap1p and Cla4p are required for regular lateral interactions between septin filaments that bring them into register to form the proposed collar. Without these factors, improper lateral interactions could cause the collapse of the collar into thick bundles of filaments, seen as bars.

Gin4p, Nap1, and Cla4p do not constitute a simple linear pathway for septin organization, because the septin defects in deletion mutants lacking any two of these proteins are significantly more severe than those observed in the single mutants [29^{**}] (Figure 2b). However, even cells lacking all three proteins do generate patches of septin-containing cortex at the neck, indicating that additional pathways exist to recruit and/or assemble septins. Recently, Bouquin *et al.* [30^{*}] showed that the protein kinase Elm1p, like Gin4p, is localized to the septin cortex and plays a role in septin organization. However, rather than displaying septin bars at the neck, cells lacking Elm1p frequently displayed ectopic septin localization near the bud tip, in addition to an irregular septin-containing zone at the neck (Figure 2b). A similar septin localization has been observed in some strains deprived of both Cla4p and the related kinase Ste20p [31,32^{*}]. Yet another pattern of septin mislocalization was seen in cells containing a particular mutant allele of *CDC42*, which frequently displayed ectopic rings of septin staining partway into the bud [33^{*}] (Figure 2b). Together, these studies indicate that many genes contribute to the normal organization of septins within the septin cortex and that a variety of curious aberrations can be produced in mutant strains. However, the basis for these aberrations and the exact role played by each protein in septin organization remain mysterious.

Interestingly, Gin4p kinase activation appears to require the septins, and it may involve recruitment of Gin4p to the neck [34]. In addition, Gin4p kinase activity is cell-cycle-regulated, with a peak during mitosis [35]. Full activation of Gin4p requires Nap1p, Cla4p and Elm1p, and these findings have been interpreted as evidence for the existence of a signaling pathway initiated by the mitotic form of the cyclin-dependent kinase Cdc28p and culminating in Gin4p activation [34–37]. However, an alternative interpretation

Figure 2

Septin organization at the neck. (a) Models of septin filament organization. Filaments have been suggested to wrap around the neck (left, [6]) or to align along the mother–bud axis (right, [25]). In the model on the left, the dark striations in Figure 1b are thought to represent individual 10 nm diameter filaments, whereas, in the model on the right, the light and dark stripes represent inherent repeat units within the filaments that are laterally associated in register with each other. (b) Aberrant septin organization in mutant strains. Note the septin bars in the *gin4Δ* and *nap1Δ* mutants, bud-tip-localized septins in the *elm1Δ* mutants, and misplaced septin rings within the buds of the *cdc42^{V36T}* mutants. Adapted, with permission, from Longtine *et al.* [29••], Bouquin *et al.* [30•] and Gladfelter *et al.* [33•].



of these results could be that the defective septins in *nap1*, *gin4* or *elm1* mutants fail to stimulate full Gin4p activation. As Gin4p kinase activity, in turn, contributes to septin organization [28], these findings suggest a bidirectional communication between septins and Gin4p that could contribute to the higher-order structure of the septin cortex. It will be very interesting to determine how septins affect Gin4p activation and whether septin organization, like Gin4p activity, is regulated during mitosis.


The septin cortex through the cell cycle

As judged either by immunofluorescence or by the use of GFP-tagged septins, a discrete ring containing five septins forms in late G1 phase, 15 minutes or so before the emergence of a visible bud (Figure 3) [34,38–42]. (Two other *S. cerevisiae* septins are expressed only during sporulation [43,44].) Following bud emergence, the ring broadens into an hourglass-shaped cortical zone at the neck, and the septins remain at the neck throughout the cell cycle. During cytokinesis, the hourglass splits into two rings that remain on both mother and daughter cells and can persist until a new ring forms in the next cell cycle. These observations raise a host of interesting questions. What causes the septin ring to form? How does the ring change into an hourglass during bud emergence? How does the septin cortex remain at the neck while growth is directed into the bud? How does the hourglass split into two rings during cytokinesis? What causes the rings that remain after cytokinesis to eventually disassemble?

Assembly of the septin ring requires the GTPase Cdc42p [42,45], which concentrates in a cortical patch at the presumptive bud site in response to cell cycle cues during late G1 [45–47]. Cdc42p is required for polarization of the actin cytoskeleton as well as formation of the septin ring, but actin polarization and septin assembly are independent

events, in that each can occur in the absence of the other [45,48]. It is not clear whether Cdc42p employs separate effector pathways to trigger these events, and the effectors linking Cdc42p to septin-ring assembly remain mysterious (although Cla4p is likely to be one of these, see above). The recent identification of *cdc42* alleles with septin-specific defects promises to accelerate progress in this area [33•].

The spreading of the septins from the ring in unbudded cells to the hourglass in budded cells may occur in response to the local cell-shape change that accompanies bud formation. Consistent with this view, cells that are prevented from forming buds by disruption of the actin cytoskeleton appear to retain septin rings for prolonged periods without spreading or other obvious changes [48]. Interestingly, the striations observed at the neck by electron microscopy were not detected before bud emergence, and did not appear fully developed until significantly after bud emergence [6,7]. Although the difficulty of visualizing these structures makes it dangerous to draw firm conclusions,

Figure 3

Septin localization through the cell cycle. (a,b) Septins initially assemble into a planar ring at the incipient bud site. (c,d) Upon bud emergence, septin staining extends into an hourglass shape at the neck. (e) At about the time of cytokinesis, the septin hourglass splits into two rings, (f) which remain, following cell separation, as an 'old' ring that is fainter and broader than the initial ring.

Figure 4




Figure 4 legend

Proteins that localize to the neck cortex. Synthesizing the available data on protein localization to the neck is complicated by the facts that the reagents and protocols employed to visualize the proteins have not been uniform and that detailed characterizations of the timing or asymmetry of localization were frequently not a primary goal of the investigators. (This figure will therefore undoubtedly contain errors, and the authors strongly encourage investigators to contact them with corrections or new information.) *To enhance the clarity of the figure, the cytokinesis remnant is indicated only in the second G1 interval, following cytokinesis. The grey bar indicates the time between disassembly of the mitotic spindle and completion of the septum.

†Timing of mRNA accumulation is drawn from the genomic study of Spellman *et al.* [62]. C, constitutively expressed. ‡Localization of these proteins to the neck has been shown to be eliminated in septin-mutant cells shifted to restrictive temperature. §Rax2p persists at old division sites even after septins and other proteins have disassembled.

¶Localization of these proteins to the neck is not eliminated in septin-mutant cells shifted to restrictive temperature. **Localization of Clb2p to the neck is only detected when its ability to concentrate in the nucleus is impaired. ††A large number of proteins localize to sites of cell growth together with Cdc42p. These include polarity establishment proteins, exocyst components and cell-wall biosynthetic enzymes. These proteins are found in a patch at the presumptive bud site in late G1, around which the septins form a ring. They then part company temporarily with the septins, localizing to the bud tip in small buds and more diffusely within larger buds, before returning to the neck at the time of cytokinesis. Letters in parentheses in the References column of the figure correspond to the following: (a) E Bi, personal communication; (b) M Longtine, K Lee, personal communication; (c) I-C Yu, JR Pringle, unpublished data; (d) A Ragnini-Wilson, personal communication; (e) L Schenkman, JR Pringle, unpublished data; (f) C Hardy, personal communication; (g) M Longtine, personal communication.

it seems likely that the transition of septins from a ring to an hourglass entails a major change in the organization of this cortical domain.

The splitting of the hourglass into two rings at the end of the cell cycle also coincides with a dramatic change in local cell shape as cells undergo cytokinesis. Interestingly, the neck striations seem to disappear before cytokinesis [6,7], and a recent study [49**] provided compelling evidence that the splitting of the septin hourglass is not simply a consequence of cell division. Mutants lacking the actomyosin ring component Iqg1p/Cyk1p are unable to undergo cytokinesis, but septin splitting occurs nevertheless. The controlling trigger for mitotic exit and cytokinesis is thought to be the GTPase Tem1p, which stimulates a signaling pathway called the ‘mitotic exit network’ to inactivate the mitotic form of Cdc28p [50,51]. Tem1p also appears to play a more direct role in cytokinesis by promoting actomyosin-ring contraction, probably through binding to Iqg1p/Cyk1p [52]. Intriguingly, it appears that Tem1p is also required for splitting of the septin hourglass, even when its role in triggering Cdc28p inactivation is bypassed [49**]. This suggests that Tem1p promotes septin splitting through a novel pathway, and it will be fascinating to learn how this works.

After cytokinesis, the septin ring remains for a time at the site of cell division. A haploid yeast cell normally forms its new bud adjacent to the site of the preceding cell division, and there is ample evidence that septins play a key role in marking the division site by recruiting several bud-site-selection proteins, such as Bud3p, Bud4p, Axl2p/Bud10p and Bud5p [53–58]. However, the old septin ring (and associated proteins) eventually disassembles, and if the cells experience a delay in entering the next cell cycle, they ‘forget’ this positional landmark [59]. It is unclear whether some specific cell-cycle signal triggers disassembly of the septin structure or if it undergoes a spontaneous decay. Some very interesting observations in this regard were made by Johnson and Blobel [60**], who demonstrated that several of the septins are covalently

modified by attachment of the ubiquitin-related protein Smt3p (or small ubiquitin-related modifier [SUMO]) (see also [61]). Smt3p was conjugated to septins during mitosis and was removed at around the time of cytokinesis (Figure 4). Remarkably, cells that were rendered largely unable to conjugate Smt3p to septins (by the heroic expedient of mapping and mutating the conjugation-site lysines in three septin genes) retained old septin rings for prolonged periods, often accumulating multiple such structures over time. This finding suggested that Smt3p influenced the eventual disassembly of the old septin structure (but see Update).

Dynamic and asymmetric character of the septin cortex

The septin cortex plays host to a multitude of arriving and departing proteins during the cell cycle (Figure 4). How is the timing of localization of individual proteins to the neck regulated? Based on genomic analyses of gene expression, 25 of the 37 non-septin proteins known to associate with the septin cortex are the products of genes that are transcribed periodically during the cell cycle [62], so periodic protein accumulation may underlie the timing of neck association. This appears to be true at least for Bud3p [23•] and Iqg1p [63], for which it has been shown that earlier synthesis of the protein during the cell cycle causes premature association with the septin cortex. However, many proteins that are present throughout the cell cycle show restricted and, in some cases, complex, timing of localization to the neck. For instance, the chitin synthase III complex (Chs3p/Chs4p), which is responsible for making the chitin ring, localizes to the septin cortex before bud emergence but departs shortly thereafter and resides in an endosomal compartment until the time of cytokinesis, when it returns to the septin cortex [15,64]. Another example is provided by Bud6p, which functions in both bud-site selection and spindle orientation [65,66]. Bud6p is initially concentrated at the presumptive bud site and the bud tip, and it subsequently begins to accumulate at the neck [65,67•]. The timing of Bud6p neck localization appears to depend on a competition between factors (including Bni1p) that attract Bud6p to the bud tip and other factors (including Bud3p) that attract

Bud6p to the neck [67^{*}]. Surprisingly, the outcome of this competition is different in haploid and diploid cells, with Bud6p accumulating at the neck much earlier in haploids. These examples illustrate the potential for cell-cycle and mating-type cues that affect membrane trafficking, polarized growth, or other processes to determine when particular proteins visit the neck.

In addition to the temporal differences in the composition of the septin cortex through the cell cycle, this membrane domain is spatially differentiated along the mother-bud axis. Although many proteins are present throughout the septin cortex, several are concentrated primarily on the mother side of the neck, some localize to a ring in the center of the neck, and others are located predominantly on the bud side of the neck (see Figure 4). Thus, the composition of the septin cortex varies along the mother–bud axis. How might this asymmetry be achieved? One appealing hypothesis was suggested on the basis of the behavior of Bni4p, which links the chitin synthase III complex to the septins [15]. Bni4p is located on the mother side of the neck during early stages of bud formation, but gradually becomes more symmetrically distributed across the neck during subsequent bud growth (Figure 4). Because of cell-cycle-regulated mRNA accumulation, the bulk of Bni4p appears to be synthesized before bud emergence, and it associates with the septin ring in unbudded cells. DeMarini and colleagues speculated that the septin cortex forming the ring before bud emergence may remain intact at the base (mother side) of the expanded hourglass cortex seen in budded cells. Thus, the pool of Bni4p made during G1 and associated with the ring at that time would remain concentrated on the mother side of the neck, and the lower levels of Bni4p synthesized later in the cell cycle may gradually populate the bud side of the septin cortex. However, more recent studies revealed that other proteins that are initially associated with the septin ring in unbudded cells do not remain on the mother side of the neck as the bud emerges. These proteins include Gin4p, which distributes throughout the neck [28], Myo1p, which localizes to the central ring [68,69], and Kcc4p, which localizes to the bud side of the neck [70]. Thus, there is no systematic correlation between the timing of recruitment to the septin cortex and localization along the mother–bud axis, suggesting that time of assembly cannot, in general, account for the asymmetry of the septin cortex.

Another hypothesis to explain the asymmetry is based on the model that septin filaments are oriented along the mother–bud axis (Figure 2a). If the filaments are intrinsically polar (like actin or tubulin polymers), and if they are all aligned in the same direction, then the septins themselves may provide an intrinsic asymmetry that could be exploited by septin-binding proteins. Thus, asymmetric localization of a septin-associated protein would simply reflect molecular recognition of the underlying septin polarity. However, this hypothesis does not easily account for proteins that localize to a ring in the middle of the neck, and it requires additional assumptions to explain the

behavior of proteins, such as Bni4p, whose distribution within the neck changes during the cell cycle.

The septin cortex as cell shape sensor

In the past few years, it has become clear that cell-cycle-checkpoint controls monitor cytoplasmic as well as nuclear events [71]. In *S. cerevisiae*, a morphogenesis checkpoint delays mitosis in response to insults that perturb bud formation [72]. This mitotic delay involves stabilization of the cell-cycle-inhibitory kinase Swe1p [73]. Targeting of Swe1p for degradation involves its interaction with two regulators, Hsl1p and Hsl7p [74]. Intriguingly, these proteins are recruited to the bud side of the septin cortex at the time of bud emergence, and their function in downregulating Swe1p appears to require this localization [29^{**},70,75]. As discussed above, the septin cortex expands from a ring to an hourglass during bud emergence, and it has been argued that Hsl1p and Hsl7p may monitor this septin reorganization and trigger Swe1p degradation only after a bud has formed [29^{**}]. In this way, the reorganization of the septin cortex in response to bud formation may allow the septins to act as indicators of local cell shape.

In fission yeast, a cytokinesis checkpoint arrests the cell cycle if cytokinesis was not completed successfully in the preceding cell cycle [76,77]. There are hints that cytokinesis may also be monitored in budding yeast. In particular, examination of mutants in which cytokinesis is slowed or prevented indicates that both the septins and cortical actin patches can persist at the neck for longer than usual under these circumstances [68,69,78^{**},79]. Does this mean that cells ‘know’ if cytokinesis has been completed, and delay the release of cytoskeletal elements from the neck when more time is required for cell division? In contrast to the mutants referred to above, septin-mutant cells proceed rapidly with the subsequent round of bud formation despite a complete failure of cytokinesis, suggesting that septins must be assembled for any ‘cytokinesis sensor’ to work. The change in local cell shape that accompanies cytokinesis may well cause alterations of septin organization analogous to those thought to accompany bud emergence. Conceivably, therefore, the septin cortex may allow cells to monitor local shape changes at both of these stages.

Conclusions

The mother–bud neck is a specialized subdomain of the cell cortex organized by the septins. Recent studies indicate that the septin cortex can function as a diffusion barrier and perhaps also as a local cell-shape sensor. To understand these functions, we must learn much more about the detailed architecture of the septin cortex and about its regulation by the cell cycle and by local geometry. The number of proteins known to visit the neck in a septin-dependent manner is large and growing, but our models of septin organization at the neck are still purely speculative. The general problem of how a cortical domain is specified and organized is of great biological importance, and we believe that investigation of septin-dependent

cortical organization will yield many insights in the coming years. Such advances will require integration of the information from molecular genetic approaches with detailed structural information. We hope that the fascinating inroads being made into septin regulation and function will attract more investigators with structural expertise into this area. The rewards should be magnificent.

Update

Recently, Johnson and Gupta [98] identified Siz1p as an E3-type enzyme responsible for conjugation of Smt3p to the septins. Like Smt3p, Siz1p was concentrated on the mother side of the neck during mitosis. A previous study [60**] found that cells in which septins had been mutated at the Smt3p-conjugation-site lysines had a defect in disassembling the old septin rings. However, cells lacking Siz1p showed no discernible defect in septin localization, suggesting that the previous result was due to an alteration of septin structure rather than to the lack of Smt3p conjugation, and the role of Smt3p at the septin cortex remains mysterious.

Acknowledgements

We thank E Bi, M Longtine, C Hardy, K Lee, J Berman and A Ragnini-Wilson for communicating results prior to publication. Thanks also go to S Kornbluth, K Bloom and members of the Lew and Pringle laboratories for many stimulating discussions. Work from the authors' laboratories was supported by National Institutes of Health grants to JRP and DJL and by funds from the Leukemia and Lymphoma Society to DJL.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Marchant R, Smith DJ: **Bud formation in *Saccharomyces cerevisiae* and a comparison with the mechanism of cell division in other yeasts.** *J Gen Microbiol* 1968, 53:163-169.
2. Hayashibe M, Katohda S: **Initiation of budding and chitin ring.** *J Gen Appl Microbiol* 1973, 19:23-39.
3. Seichertová O, Beran K, Holan Z, Pokorný V: **The chitin-glucan complex of *Saccharomyces cerevisiae*. II. Location of the complex in the encircling region of the bud scar.** *Folia Microbiol* 1973, 18:207-211.
4. Cabib E, Roh DH, Schmidt M, Crotti LB, Varma A: **The yeast cell wall and septum as paradigms of cell growth and morphogenesis.** *J Biol Chem* 2001, 276:19679-19682.
5. Longtine MS, DeMarini DJ, Valencik ML, Al-Awar OS, Fares H, De Virgilio C, Pringle JR: **The septins: roles in cytokinesis and other processes.** *Curr Opin Cell Biol* 1996, 8:106-119.
6. Byers B, Goetsch L: **A highly ordered ring of membrane-associated filaments in budding yeast.** *J Cell Biol* 1976, 69:717-721.
7. Byers B: **Cytology of the yeast life cycle.** In *The Molecular Biology of the Yeast *Saccharomyces: Life Cycle and Inheritance**. Edited by Strathern JN, Jones EW, Broach JR. Cold Spring Harbor Laboratory; 1981:59-96.
8. Neufeld TP, Rubin GM: **The *Drosophila* peanut gene is required for cytokinesis and encodes a protein similar to yeast putative bud neck filament proteins.** *Cell* 1994, 77:371-379.
9. Fares H, Peifer M, Pringle JR: **Localization and possible functions of *Drosophila* septins.** *Mol Biol Cell* 1995, 6:1843-1859.
10. Kinoshita M, Kumar S, Mizoguchi A, Ide C, Kinoshita A, Haraguchi T, Hiraoka Y, Noda M: **Nedd5, a mammalian septin, is a novel cytoskeletal component interacting with actin-based structures.** *Genes Dev* 1997, 11:1535-1547.
11. Adam JC, Pringle JR, Peifer M: **Evidence for functional differentiation among *Drosophila* septins in cytokinesis and cellularization.** *Mol Biol Cell* 2000, 11:3123-3135.
12. Nguyen TQ, Sawa H, Okano H, White JG: **The *C. elegans* septin genes, *unc-59* and *unc-61*, are required for normal postembryonic cytokinesis and morphogenesis but have no essential function in embryogenesis.** *J Cell Sci* 2000, 113:3825-3837.
13. Kartmann B, Roth D: **Novel roles for mammalian septins: from vesicle trafficking to oncogenesis.** *J Cell Sci* 2001, 114:839-844.
14. Trimble WS: **Septins: a highly conserved family of membrane-associated GTPases with functions in cell division and beyond.** *J Membr Biol* 1999, 169:75-81.
15. DeMarini DJ, Adams AEM, Fares H, De Virgilio C, Valle G, Chuang JS, Pringle JR: **A septin-based hierarchy of proteins required for localized deposition of chitin in the *Saccharomyces cerevisiae* cell wall.** *J Cell Biol* 1997, 139:75-93.
16. Chant J: **Cell polarity in yeast.** *Annu Rev Cell Dev Biol* 1999, 15:365-391.
17. Hales KG, Bi E, Wu JQ, Adam JC, Yu IC, Pringle JR: **Cytokinesis: an emerging unified theory for eukaryotes?** *Curr Opin Cell Biol* 1999, 11:717-725.
18. Takizawa PA, DeRisi JL, Wilhelm JE, Vale RD: **Plasma membrane compartmentalization in yeast by messenger RNA transport and a septin diffusion barrier.** *Science* 2000, 290:341-344.
- The asymmetric localization of the membrane protein Ist2p to the bud, but not the mother, plasma membrane is shown to be due to a septin-dependent diffusion barrier at the neck.
19. Barral Y, Mermall V, Mooseker MS, Snyder M: **Compartmentalization of the cell cortex by septins is required for maintenance of cell polarity in yeast.** *Mol Cell* 2000, 5:841-851.
- Bud growth occurs in two stages. In cells with small buds, new cell wall is deposited mainly at the bud tip (apical growth) and, in cells with larger buds, new cell wall is deposited all over the bud surface (isotropic growth). Septins are shown to be important for maintaining the bud-specific localization of several peripheral membrane proteins specifically during isotropic bud growth. In addition, filming of cortical actin patch movements in living cells shows that individual patches can cross from the bud into the mother. However, patches in the mother are relatively short-lived, accounting (at least in part) for the concentration of actin patches within the bud.
20. Doyle T, Botstein D: **Movement of yeast cortical actin cytoskeleton visualized *in vivo*.** *Proc Natl Acad Sci USA* 1996, 93:3886-3891.
21. Waddle JA, Karpova TS, Waterston RH, Cooper JA: **Movement of cortical actin patches in yeast.** *J Cell Biol* 1996, 132:861-870.
22. Pelham RJ Jr, Chang F: **Role of actin polymerization and actin cables in actin-patch movement in *Schizosaccharomyces pombe*.** *Nat Cell Biol* 2001, 3:235-244.
23. Lord M, Yang MC, Mischke M, Chant J: **Cell cycle programs of gene expression control morphogenetic protein localization.** *J Cell Biol* 2000, 151:1501-1511.
- Periodic mRNA accumulation during the cell cycle programs synthesis of Bud10p during late G1, and synthesis of Bud3p during G2/M. Both proteins associate with the neck cortex in the second half of the cell cycle. Early or constitutive synthesis of Bud3p results in early recruitment of Bud3p to the septin cortex, but has no detrimental consequences for Bud3p function. In contrast, constitutive or late synthesis of the membrane protein Bud10p results in localization of Bud10p throughout the plasma membrane, causing a significant impairment of Bud10p function. The association of Bud10p with the neck normally occurs after its delivery to the plasma membrane, and persists even beyond cytokinesis, so it is quite surprising that delayed arrival of Bud10p prevents its effective localization to the neck.
24. Gale C, Gerami-Nejad M, McClellan M, Vandoninck S, Longtine MS, Berman J: ***Candida albicans* Int1p interacts with the septin ring in yeast and hyphal cells.** *Mol Biol Cell* 2001, 12:in press.
25. Field CM, Al-Awar O, Rosenblatt J, Wong ML, Alberts B, Mitchison TJ: **A purified *Drosophila* septin complex forms filaments and exhibits GTPase activity.** *J Cell Biol* 1996, 133:605-616.
26. Frazier JA, Wong ML, Longtine MS, Pringle JR, Mann M, Mitchison TJ, Field C: **Polymerization of purified yeast septins: evidence that organized filament arrays may not be required for septin function.** *J Cell Biol* 1998, 143:737-749.
27. Field CM, Kellogg D: **Septins: cytoskeletal polymers or signalling GTPases?** *Trends Cell Biol* 1999, 9:387-394.
28. Longtine MS, Fares H, Pringle JR: **Role of the yeast Gin4p protein kinase in septin assembly and the relationship between septin assembly and septin function.** *J Cell Biol* 1998, 143:719-736.

29. Longtine MS, Theesfeld CL, McMillan JN, Weaver E, Pringle JR, ●● Lew DJ: **Septin-dependent assembly of a cell-cycle-regulatory module in *Saccharomyces cerevisiae***. *Mol Cell Biol* 2000, 20:4049-4061.
- Like *gin4* mutants (see [28]), *cla4* and *nap1* mutants display defects in septin organization, with septins frequently forming bars at the neck. Septin misorganization is more severe in double mutants, and these mutants also have very elongated buds that arise because of a Swe1p-dependent delay in the cell cycle (see also [70]). Two negative regulators of Swe1p, Hsl1p and Hsl7p, localize to the bud side of the septin cortex in budded cells and recruit Swe1p to that site. However, Hsl1p and Hsl7p are released from the neck in mutants with defective septins. The authors suggest that recruitment of these proteins to the septin cortex is required for Swe1p downregulation, and that this recruitment may depend on the septin reorganization that occurs during bud formation.
30. Bouquin N, Barral Y, Courbeyrette R, Blondel M, Snyder M, Mann C: ●● **Regulation of cytokinesis by the Elm1 protein kinase in *Saccharomyces cerevisiae***. *J Cell Sci* 2000, 113:1435-1445.
- The Elm1p protein kinase localizes to the septin cortex and *elm1* mutants display elongated buds with septins mislocalized to the bud tip.
31. Cvrcková F, De Virgilio C, Manser E, Pringle JR, Nasmyth K: **Ste20-like protein kinases are required for normal localization of cell growth and for cytokinesis in budding yeast**. *Genes Dev* 1995, 9:1817-1830.
32. Weiss EL, Bishop AC, Shokat KM, Drubin DG: ●● **Chemical genetic analysis of the budding-yeast p21-activated kinase Cla4p**. *Nat Cell Biol* 2000, 2:677-685.
- Using a *cla4* allele that encodes a product that can be reversibly inactivated by a specific inhibitor, the authors investigate the roles of the related Cla4p and Ste20p kinases. When kinase activity is inhibited at the time of septin-ring formation, septins localize aberrantly to bud tips and cells make elongated buds. If the inhibitor is then removed, a new septin ring forms at the bud tip. Interestingly, Cla4p/Ste20p inhibition has no effect once a septin ring has formed.
33. Gladfelter AS, Moskow JJ, Zyla TR, Lew DJ: ●● **Isolation and characterization of effector-loop mutants of *CDC42* in yeast**. *Mol Biol Cell* 2001, 12:1239-1255.
- An analysis of *cdc42* mutants uncovers two alleles that display mislocalized septin rings (frequently midway into the bud) and have unusually wide necks.
34. Carroll CW, Altman R, Schieltz D, Yates JR, Kellogg D: **The septins are required for the mitosis-specific activation of the Gin4 kinase**. *J Cell Biol* 1998, 143:709-717.
35. Altman R, Kellogg D: **Control of mitotic events by Nap1 and the Gin4 kinase**. *J Cell Biol* 1997, 138:119-130.
36. Sreenivasan A, Kellogg D: **The Elm1 kinase functions in a mitotic signaling network in budding yeast**. *Mol Cell Biol* 1999, 19:7983-7994.
37. Tjandra H, Compton J, Kellogg D: **Control of mitotic events by the Cdc42 GTPase, the Clb2 cyclin and a member of the PAK kinase family**. *Curr Biol* 1998, 8:991-1000.
38. Haarer BK, Pringle JR: **Immunofluorescence localization of the *Saccharomyces cerevisiae* CDC12 gene product to the vicinity of the 10-nm filaments in the mother-bud neck**. *Mol Cell Biol* 1987, 7:3678-3687.
39. Kim HB, Haarer BK, Pringle JR: **Cellular morphogenesis in the *Saccharomyces cerevisiae* cell cycle: localization of the CDC3 gene product and the timing of events at the budding site**. *J Cell Biol* 1991, 112:535-544.
40. Ford SK, Pringle JR: **Cellular morphogenesis in the *Saccharomyces cerevisiae* cell cycle: localization of the CDC11 gene product and the timing of events at the budding site**. *Dev Genet* 1991, 12:281-292.
41. Mino A, Tanaka K, Kamei T, Umikawa M, Fujiwara T, Takai Y: **Shs1p: a novel member of septin that interacts with spa2p, involved in polarized growth in *Saccharomyces cerevisiae***. *Biochem Biophys Res Commun* 1998, 251:732-736.
42. Cid VJ, Adamikova L, Sanchez M, Molina M, Nombela C: **Cell cycle control of septin ring dynamics in the budding yeast**. *Microbiology* 2001, 147:1437-1450.
43. Fares H, Goetsch L, Pringle JR: **Identification of a developmentally regulated septin and involvement of the septins in spore formation in *Saccharomyces cerevisiae***. *J Cell Biol* 1996, 132:399-411.
44. De Virgilio C, DeMarini DJ, Pringle JR: **SPR28, a sixth member of the septin gene family in *Saccharomyces cerevisiae* that is expressed specifically in sporulating cells**. *Microbiology* 1996, 142:2897-2905.
45. Pringle JR, Bi E, Harkins HA, Zahner JE, De Virgilio C, Chant J, Corrado K, Fares H: **Establishment of cell polarity in yeast**. *Cold Spring Harbor Symp Quant Biol* 1995, 60:729-744.
46. Ziman M, Preuss D, Mulholland J, O'Brien JM, Botstein D, Johnson DI: **Subcellular localization of Cdc42p, a *Saccharomyces cerevisiae* GTP-binding protein involved in the control of cell polarity**. *Mol Biol Cell* 1993, 4:1307-1316.
47. Gulli MP, Jaquenoud M, Shimada Y, Niederhauser G, Wiget P, Peter M: **Phosphorylation of the Cdc42 exchange factor Cdc24 by the PAK-like kinase Cla4 may regulate polarized growth in yeast**. *Mol Cell* 2000, 6:1155-1167.
48. Ayscough KR, Stryker J, Pokala N, Sanders M, Crews P, Drubin DG: **High rates of actin filament turnover in budding yeast and roles for actin in establishment and maintenance of cell polarity revealed using the actin inhibitor latrunculin-A**. *J Cell Biol* 1997, 137:399-416.
49. Lippincott J, Shannon KB, Shou W, Deshaies RJ, Li R: ●● **The Tem1 small GTPase controls actomyosin and septin dynamics during cytokinesis**. *J Cell Sci* 2001, 114:1379-1386.
- The small GTPase Tem1p controls mitotic exit via a signaling pathway that releases the phosphatase Cdc14p from the nucleolus, allowing it to inactivate mitotic CDK complexes. The authors now show that Tem1p is required for the splitting of the septin hourglass structure into two rings at the time of cytokinesis, even in mutant strains in which Cdc14p is constitutively released from the nucleolus. Tem1p is also required for actomyosin ring contraction, but neither ring contraction nor cytokinesis *per se* are required for septin splitting. The authors suggest that Tem1p-dependent septin splitting may facilitate actomyosin ring contraction, and thereby promote cytokinesis.
50. Morgan DO: **Regulation of the APC and the exit from mitosis**. *Nat Cell Biol* 1999, 1:E47-E53.
51. McCollum D, Gould KL: **Timing is everything: regulation of mitotic exit and cytokinesis by the MEN and SIN**. *Trends Cell Biol* 2001, 11:89-95.
52. Shannon KB, Li R: **The multiple roles of Cyk1p in the assembly and function of the actomyosin ring in budding yeast**. *Mol Biol Cell* 1999, 10:283-296.
53. Chant J, Mischke M, Mitchell E, Herskowitz I, Pringle JR: **Role of Bud3p in producing the axial budding pattern of yeast**. *J Cell Biol* 1995, 129:767-778.
54. Sanders SL, Herskowitz I: **The Bud4 protein of yeast, required for axial budding, is localized to the mother/bud neck in a cell cycle-dependent manner**. *J Cell Biol* 1996, 134:413-427.
55. Halme A, Michelitch M, Mitchell EL, Chant J: **Bud10p directs axial cell polarization in budding yeast and resembles a transmembrane receptor**. *Curr Biol* 1996, 6:570-579.
56. Marston AL, Chen T, Yang MC, Belhumeur P, Chant J: **A localized GTPase exchange factor, Bud5, determines the orientation of division axes in yeast**. *Curr Biol* 2001, 11:803-807.
57. Roemer T, Madden K, Chang J, Snyder M: **Selection of axial growth sites in yeast requires Axl2p, a novel plasma membrane glycoprotein**. *Genes Dev* 1996, 10:777-793.
58. Kang PJ, Sanson A, Lee B, Park HO: **A GDP/GTP exchange factor involved in linking a spatial landmark to cell polarity**. *Science* 2001, 292:1376-1378.
59. Chant J, Pringle JR: **Patterns of bud-site selection in the yeast *Saccharomyces cerevisiae***. *J Cell Biol* 1995, 129:751-765.
60. Johnson ES, Blobel G: ●● **Cell cycle-regulated attachment of the ubiquitin-related protein SUMO to the yeast septins**. *J Cell Biol* 1999, 147:981-993.
- This paper documents the cell-cycle-dependent modification of three septins by the SUMO homolog, Smt3p. Inability to conjugate Smt3p causes the persistence of septin rings for a long time after completion of cytokinesis.
61. Takahashi Y, Iwase M, Konishi M, Tanaka M, Toh-e A, Kikuchi Y: **Smt3, a SUMO-1 homolog, is conjugated to Cdc3, a component of septin rings at the mother-bud neck in budding yeast**. *Biochem Biophys Res Commun* 1999, 259:582-587.
62. Spellman PT, Sherlock G, Zhang MQ, Iyer VR, Anders K, Eisen MB, Brown PO, Botstein D, Futcher B: **Comprehensive identification of cell cycle-regulated genes of the yeast *Saccharomyces cerevisiae* by microarray hybridization**. *Mol Biol Cell* 1998, 9:3273-3297.
63. Epp JA, Chant J: **An IQGAP-related protein controls actin-ring formation and cytokinesis in yeast**. *Curr Biol* 1997, 7:921-929.

64. Chuang JS, Schekman RW: Differential trafficking and timed localization of two chitin synthase proteins, Chs2p and Chs3p. *J Cell Biol* 1996, 135:597-610.
65. Amberg DC, Zahner JE, Mulholland JW, Pringle JR, Botstein D: Aip3p/Bud6p, a yeast actin-interacting protein that is involved in morphogenesis and the selection of bipolar budding sites. *Mol Biol Cell* 1997, 8:729-753.
66. Yeh E, Yang C, Chin E, Maddox P, Salmon ED, Lew DJ, Bloom K: Dynamic positioning of mitotic spindles in yeast: role of microtubule motors and cortical determinants. *Mol Biol Cell* 2000, 11:3949-3961.
67. Segal M, Bloom K, Reed SI: Bud6 directs sequential microtubule interactions with the bud tip and bud neck during spindle morphogenesis in *Saccharomyces cerevisiae*. *Mol Biol Cell* 2000, 11:3689-3702.
- A very detailed analysis of Bud6p localization in haploid and diploid cells, as well as various mutants, is presented. Previous results had indicated that certain cell-cycle perturbations caused a lethal phenotype that involved irreversible migration of the mitotic spindle into the bud in diploid, but not in haploid, cells. The differences in Bud6p behaviour explain these perplexing observations and suggest that Bud6p is an important cortical cue used during orientation of the spindle along the mother–bud axis.
68. Bi E, Maddox P, Lew DJ, Salmon ED, McMillan JN, Yeh E, Pringle JR: Involvement of an actomyosin contractile ring in *Saccharomyces cerevisiae* cytokinesis. *J Cell Biol* 1998, 142:1301-1312.
69. Lippincott J, Li R: Sequential assembly of myosin II, an IQGAP-like protein, and filamentous actin to a ring structure involved in budding yeast cytokinesis. *J Cell Biol* 1998, 140:355-366.
70. Barral Y, Parra M, Bidlingmaier S, Snyder M: Nim1-related kinases coordinate cell cycle progression with the organization of the peripheral cytoskeleton in yeast. *Genes Dev* 1999, 13:176-187.
71. Lew DJ: Cell-cycle checkpoints that ensure coordination between nuclear and cytoplasmic events in *Saccharomyces cerevisiae*. *Curr Opin Genet Dev* 2000, 10:47-53.
72. Lew DJ, Reed SI: A cell cycle checkpoint monitors cell morphogenesis in budding yeast. *J Cell Biol* 1995, 129:739-749.
73. Sia RAL, Bardes ESG, Lew DJ: Control of Swe1p degradation by the morphogenesis checkpoint. *EMBO J* 1998, 17:6678-6688.
74. McMillan JN, Longtine MS, Sia RAL, Theesfeld CL, Bardes ESG, Pringle JR, Lew DJ: The morphogenesis checkpoint in *Saccharomyces cerevisiae*: cell cycle control of Swe1p degradation by Hsl1p and Hsl7p. *Mol Cell Biol* 1999, 19:6929-6939.
75. Shulewitz MJ, Inouye CJ, Thorner J: Hsl7 localizes to a septin ring and serves as an adapter in a regulatory pathway that relieves tyrosine phosphorylation of Cdc28 protein kinase in *Saccharomyces cerevisiae*. *Mol Cell Biol* 1999, 19:7123-7137.
76. Le Goff X, Woollard A, Simanis V: Analysis of the *cps1* gene provides evidence for a septation checkpoint in *Schizosaccharomyces pombe*. *Mol Gen Genet* 1999, 262:163-172.
77. Liu J, Wang H, Balasubramanian MK: A checkpoint that monitors cytokinesis in *Schizosaccharomyces pombe*. *J Cell Sci* 2000, 113:1223-1230.
78. Vallen EA, Caviston J, Bi E: Roles of Hof1p, Bni1p, Bnr1p, and Myo1p in cytokinesis in *Saccharomyces cerevisiae*. *Mol Biol Cell* 2000, 11:593-611.
- This detailed study examines the roles of several neck-localized proteins in cytokinesis. The authors argue that cytokinesis can proceed through either of two mechanistically distinct 'pathways' – one involving contraction of the actomyosin ring and the other involving deposition of septal material at the neck. These pathways are normally coupled to each other and are both dependent on septins. The localization of Hof1p at the neck is particularly complex, exhibiting four different patterns as cells traverse the cell cycle.
79. Lippincott J, Li R: Dual function of Cyk2, a cdc15/PSTPIP family protein, in regulating actomyosin ring dynamics and septin distribution. *J Cell Biol* 1998, 143:1947-1960.
80. Moriya H, Isono K: Analysis of genetic interactions between DHH1, SSD1 and ELM1 indicates their involvement in cellular morphology determination in *Saccharomyces cerevisiae*. *Yeast* 1999, 15:481-496.
81. Boyne JR, Yosuf HM, Bieganowski P, Brenner C, Price C: Yeast myosin light chain, Mlc1p, interacts with both IQGAP and class II myosin to effect cytokinesis. *J Cell Sci* 2000, 113:4533-4543.
82. Kamei T, Tanaka K, Hihara T, Umikawa M, Imamura H, Kikyo M, Ozaki K, Takai Y: Interaction of Bnr1p with a novel Src homology 3 domain-containing Hof1p. Implication in cytokinesis in *Saccharomyces cerevisiae*. *J Biol Chem* 1998, 273:28341-28345.
83. Korinek WS, Bi E, Epp JA, Wang L, Ho J, Chant J: Cyk3, a novel SH3-domain protein, affects cytokinesis in yeast. *Curr Biol* 2000, 10:947-950.
84. Park HO, Sanson A, Herskowitz I: Localization of Bud2p, a GTPase-activating protein necessary for programming cell polarity in yeast to the presumptive bud site. *Genes Dev* 1999, 13:1912-1917.
85. Harkins HA, Page N, Schenkman L, De Virgilio C, Shaw S, Bussey H, Pringle JR: Bud8p and Bud9p, proteins that may mark the sites for bipolar budding in yeast. *Mol Biol Cell* 2001, 12:2497-2518.
86. Chen T, Hiroko T, Chaudhuri A, Inose F, Lord M, Tanaka S, Chant J, Fujita A: Multigenerational cortical inheritance of the Rax2 protein in orienting polarity and division in yeast. *Science* 2000, 290:1975-1978.
87. Hood JK, Hwang WW, Silver PA: The *Saccharomyces cerevisiae* cyclin Clb2p is targeted to multiple subcellular locations by cis- and trans-acting determinants. *J Cell Sci* 2001, 114:589-597.
88. Song S, Grenfell TZ, Garfield S, Erikson RL, Lee KS: Essential function of the polo box of Cdc5 in subcellular localization and induction of cytokinetic structures. *Mol Cell Biol* 2000, 20:286-298.
89. Frenz LM, Lee SE, Fesquet D, Johnston LH: The budding yeast Dbf2 protein kinase localises to the centrosome and moves to the bud neck in late mitosis. *J Cell Sci* 2000, 113:3399-3408.
90. Yoshida S, Toh-e A: Regulation of the localization of Dbf2 and Mob1 during cell division of *Saccharomyces cerevisiae*. *Genes Genet Syst* 2001, 76:141-147.
91. Brizzio V, Gammie AE, Rose MD: Rvs161p interacts with Fus2p to promote cell fusion in *Saccharomyces cerevisiae*. *J Cell Biol* 1998, 141:567-584.
92. Reiser V, Salah SM, Ammerer G: Polarized localization of yeast Pbs2 depends on osmolarity, the membrane protein Sho1 and Cdc42. *Nat Cell Biol* 2000, 2:620-627.
93. Marcoux N, Cloutier S, Zakrzewska E, Charest PM, Bourbonnais Y, Pallotta D: Suppression of the profilin-deficient phenotype by the RHO2 signaling pathway in *Saccharomyces cerevisiae*. *Genetics* 2000, 156:579-592.
94. Robinson LC, Bradley C, Bryan JD, Jerome A, Kweon Y, Panek HR: The Yck2 yeast casein kinase 1 isoform shows cell cycle-specific localization to sites of polarized growth and is required for proper septin organization. *Mol Biol Cell* 1999, 10:1077-1092.
95. Bloecher A, Tatchell K: Dynamic localization of protein phosphatase type 1 in the mitotic cell cycle of *Saccharomyces cerevisiae*. *J Cell Biol* 2000, 149:125-140.
96. Pruyne D, Bretscher A: Polarization of cell growth in yeast. I. Establishment and maintenance of polarity states. *J Cell Sci* 2000, 113:365-375.
97. Pruyne D, Bretscher A: Polarization of cell growth in yeast. *J Cell Sci* 2000, 113:571-585.
98. Johnson ES, Gupta AA: An E3-like factor that promotes SUMO conjugation to the yeast septins. *Cell* 2001, 106:735-744.